

# Conjectures that should be true\*

August 24, 2016

## 1 Edges in list-critical graphs

A graph  $G$  is *k-list-critical* if  $G$  is not  $(k - 1)$ -choosable, but every proper subgraph of  $G$  is  $(k - 1)$ -choosable. Replace ‘ $(k - 1)$ -’ with ‘online  $(k - 1)$ -’ and ‘ $k$ -’ with ‘*online k*-’ in the previous sentence and read it.

**Conjecture 1.** *Every incomplete k-list-critical graph has average degree at least*

$$k - 1 + \frac{k - 3}{(k - 1)^2}.$$

**Background.** The connected graphs in which each block is a complete graph or an odd cycle are called *Gallai trees*. Gallai [11] proved that in a  $k$ -critical graph, the vertices of degree  $k - 1$  induce a disjoint union of Gallai trees. The same is true for  $k$ -list-critical graphs [1, 10]. This quickly implies a lower bound on the average degree of  $k$ -list-critical graphs of

$$k - 1 + \frac{k - 3}{k^2 - 3}.$$

In [21], R. improved this to

$$k - 1 + \frac{k - 3}{k^2 - 2k + 2}$$

using a lemma from Kierstead and R. [14] that generalizes a kernel technique of Kostochka and Yancey [15]. As noted at the end of [21], a small improvement to the argument would yield Conjecture 1. [This is now known to hold for  \$k \geq 6\$ , only the  \$k = 4\$  and  \$k = 5\$  cases remain.](#) □

**Conjecture 2.** *Every incomplete online k-list-critical graph  $G$  has*

$$2 \|G\| \geq (k - 1) |G| + k - 3.$$

**Background.** Dirac [9] proved this for  $k$ -critical graphs. Kostochka and Stiebitz [16] proved it for  $k$ -list-critical graphs. Their proof does not seem to generalize. When  $|G|$  is large compared with  $k$ , the conjecture holds by Gallai-type bounds on the average degree of online  $k$ -list-critical graphs [13, 2]. □

---

\*(dis)proofs  $\Rightarrow$  [landon.rabern@gmail.com](mailto:landon.rabern@gmail.com)

## 1.1 The $\frac{5}{6}$ bound

**Conjecture 3.** *Every vertex-transitive graph has  $\chi \leq \max \left\{ \omega, \left\lceil \frac{5\Delta+3}{6} \right\rceil \right\}$ .*

**Background.** In [3], the following was proved

- the conjecture holds for the fractional chromatic number  $\chi^*$ ,
- the conjecture holds with  $\left\lceil \frac{5\Delta+3}{6} \right\rceil$  replaced by  $\epsilon(\Delta + 1)$  for some  $\epsilon < 1$ ,
- the conjecture holds both if Reed's  $\omega, \Delta, \chi$  conjecture and the strong  $2\Delta$ -colorability conjecture hold for vertex-transitive graphs (only strong  $\frac{5}{2}\Delta$ -colorability is required),

Does the conjecture hold for Cayley graphs? □

**Conjecture 4.** *Every line graph (of a multigraph) has  $\chi \leq \max \left\{ \omega, \left\lceil \frac{5\Delta+3}{6} \right\rceil \right\}$ .*

**Background.** In [18] this was proved with  $\left\lceil \frac{5\Delta+3}{6} \right\rceil$  replaced by  $\frac{7\Delta+10}{8}$ . Conjecture 14 in [18] that implies this conjecture is now known to be false. [This conjecture is true, recently proved with Dan Cranston.](#) □

## 2 Around Planar graphs

**Conjecture 5.** *Every graph with no  $K_5$ -subdivision is 2-fold 9-colorable.*

**Background.** In [4], Cranston and R. gave a short proof of this conjecture with  $K_5$ -minors excluded instead of  $K_5$ -subdivisions (which also follows from the Four Color Theorem). Hajós conjectured that every graph is  $(k - 1)$ -colorable unless it contains a subdivision of  $K_k$ . This is known to be true for  $k \leq 4$  and false for  $k \geq 7$ . The cases  $k = 5$  and  $k = 6$  remain unresolved. □

## 3 Maximum degree, clique number and colorings

### 3.1 Around Borodin-Kostochka

**Conjecture 6.** *Every graph with  $\chi \geq \Delta \geq 8$  contains a  $K_3 \vee H$  where  $H$  is some graph on  $\Delta - 3$  vertices.*

**Background.** By results in [8], for  $\Delta \geq 9$  the existence of  $K_3 \vee H$  implies the existence of  $K_\Delta$ . So, this (seemingly weaker) conjecture for  $\Delta \geq 9$  implies the Borodin-Kostochka conjecture. The one known connected counterexample to the Borodin-Kostochka conjecture for  $\Delta = 8$  is a 5-cycle with each vertex blown up to a triangle. This graph is not a counterexample to Conjecture 6. □

**Conjecture 7.** *Every graph with  $\chi \geq \Delta$  contains  $K_{\Delta-3}$ .*

**Background.** Results in [5] show that this holds with  $K_{\Delta-4}$  instead of  $K_{\Delta-3}$ . Moreover, [5] proves the conjecture for all but  $\Delta \in \{6, 8, 9, 11, 12\}$ . Reed's conjecture [22] that every graph satisfies  $\chi \leq \left\lceil \frac{\omega + \Delta + 1}{2} \right\rceil$  implies this conjecture with  $K_{\Delta-2}$  instead of  $K_{\Delta-3}$ . □

**Conjecture 8.** *Every graph with  $\chi \geq \Delta$  either contains  $K_\Delta$  or contains a  $K_{\Delta-4}$  with all  $\Delta$ -vertices.*

**Background.** Results in [5] show that this holds with  $K_{\Delta-5}$  instead of  $K_{\Delta-4}$ . For  $\Delta \leq 7$ , the conjecture holds by [19, 17]. Also by [5], it holds when  $\Delta = 3r + 1$  for  $r \geq 3$ .  $\square$

**Conjecture 9.** *Every graph with  $\Delta \geq 8$  and  $\omega < \Delta$  is 2-fold  $(2\Delta - 1)$ -colorable.*

**Background.** The one known connected counterexample to the Borodin-Kostochka conjecture for  $\Delta = 8$  is a 5-cycle with each vertex blown up to a triangle. This graph is not a counterexample to Conjecture 9.  $\square$

**Conjecture 10.** *Every graph with  $\theta \geq 10$  and  $\omega \leq \frac{\theta}{2}$  is  $\lfloor \frac{\theta}{2} \rfloor$ -choosable.*

**Background.** Here  $\theta$  is the Ore degree given by  $\theta(G) := \max_{xy \in E(G)} d(x) + d(y)$ . This conjecture holds for ordinary coloring [12, 19, 17, 20]. In [14], the conjecture is proved for  $\theta \geq 18$  for both list-coloring and online list-coloring. Further lowering of  $\theta$  would follow from improved bounds on average degree of list-critical graphs [13].  $\square$

**Conjecture 11.** *Every claw-free graph with  $\Delta \geq 9$  and  $\omega < \Delta$  is  $(\Delta - 1)$ -choosable.*

**Background.** In [7], this was proved for ordinary coloring. In [6], the conjecture was proved for  $\Delta \geq 69$ . Also, [6] proved that the full conjecture follows from the line-graph case.  $\square$

**Conjecture 12.** *There is a polynomial time graph algorithm that finds either a  $(\Delta - 1)$ -coloring or a  $K_{\Delta-3}$ .*

**Background.** In [5], the following was proved

- the conjecture holds with  $K_{\Delta-3}$  replaced by  $K_{\Delta-4}$ ,
- the conjecture holds for  $\Delta \geq 25$  (the proof uses algorithmic versions of the local lemma),
- the conjecture holds when  $\Delta = 3r + 1$ .

$\square$

## References

- [1] O.V. Borodin, *Criterion of chromaticity of a degree prescription*, Abstracts of IV All-Union Conf. on Th. Cybernetics, 1977, pp. 127–128.
- [2] D. Cranston and L. Rabern, *Edge lower bounds for list critical graphs, via discharging*, arXiv:1602.02589 (2016).
- [3] Daniel W. Cranston and Landon Rabern, *A note on coloring vertex-transitive graphs*, arXiv:1404.6550 (2014).

- [4] ———, *Planar graphs are  $9/2$ -colorable and have independence ratio at least  $3/13$* , arXiv:1410.7233 (2014).
- [5] ———, *Graphs with  $\chi=\Delta$  Have Big Cliques*, SIAM Journal on Discrete Mathematics **29** (2015), no. 4, 1792–1814.
- [6] ———, *List-coloring claw-free graphs with  $\Delta - 1$  colors*, arXiv : 1508.03574(2015).
- [7] D.W. Cranston and L. Rabern, *Coloring claw-free graphs with  $\Delta - 1$  colors*, Arxiv preprint arXiv:1206.1269 (2012).
- [8] ———, *Conjectures equivalent to the Borodin-Kostochka Conjecture that appear weaker*, Arxiv preprint arXiv:1203.5380 (2012).
- [9] G.A. Dirac, *A theorem of R.L. Brooks and a conjecture of H. Hadwiger*, *Proceedings of the London Mathematical Society* **3** (1957), no. 1, 161–195.
- [10] P. Erdős, A.L. Rubin, and H. Taylor, *Choosability in graphs*, *Proc. West Coast Conf. on Combinatorics, Graph Theory and Computing, Congressus Numerantium, vol. 26, 1979*, pp. 125–157.
- [11] T. Gallai, *Kritische Graphen I.*, *Publ. Math. Inst. Hungar. Acad. Sci* **8** (1963), 165–192 (in German).
- [12] H.A. Kierstead and A.V. Kostochka, *Ore-type versions of Brooks’ theorem*, *Journal of Combinatorial Theory, Series B* **99** (2009), no. 2, 298–305.
- [13] H.A. Kierstead and L. Rabern, *Improved lower bounds on the number of edges in list critical and online list critical graphs*, arXiv preprint arXiv:1406.7355 (2014).
- [14] ———, *Extracting list colorings from large independent sets*, arXiv:1512.08130 (2015).
- [15] Alexandr Kostochka and Matthew Yancey, *Ore’s conjecture on color-critical graphs is almost true*, *J. Combin. Theory Ser. B* **109** (2014), 73–101. MR 3269903
- [16] Alexandr V Kostochka and Michael Stiebitz, *A list version of dirac’s theorem on the number of edges in colour-critical graphs*, *Journal of Graph Theory* **39** (2002), no. 3, 165–177.
- [17] A.V. Kostochka, L. Rabern, and M. Stiebitz, *Graphs with chromatic number close to maximum degree*, *Discrete Mathematics* **312** (2012), no. 6, 1273–1281.
- [18] L. Rabern, *A strengthening of Brooks’ Theorem for line graphs*, *Electron. J. Combin.* **18** (2011), no. p145, 1.
- [19] ———,  *$\Delta$ -critical graphs with small high vertex cliques*, *Journal of Combinatorial Theory, Series B* **102** (2012), no. 1, 126–130.
- [20] ———, *Partitioning and coloring graphs with degree constraints*, *Discrete Mathematics* **9** (2013), no. 313, 1028–1034.

[21] Landon Rabern, A better lower bound on average degree of 4-list-critical graphs, *arXiv:1602.08532* (2016).

[22] B. Reed,  $\omega$ ,  $\Delta$ , and  $\chi$ , *Journal of Graph Theory* **27** (1998), no. 4, 177–212.