

A SIMILAR SHORTER PROOF OF BROOKS' THEOREM

LANDON RABERN

ABSTRACT. Lovász gave a short proof of Brooks' theorem by coloring greedily in a good order. We give a different short proof by reducing to the cubic case. Then we show how to extend the result to (online) list coloring via the Kernel Lemma.

In [7] Lovász gave a short proof of Brooks' Theorem [1] by coloring greedily in a good order. Here we give a different short proof by reducing to the cubic case. One interesting feature of the proof is that it doesn't use any connectivity concepts. Our notation follows Diestel [2] except we write K_t instead of K^t for the complete graph on t vertices.

Brooks' Theorem. *Every graph satisfies $\chi \leq \max\{3, \omega, \Delta\}$.*

Proof. Suppose the theorem is false and choose a counterexample G minimizing $|G|$. Put $\Delta := \Delta(G)$. Using minimality of $|G|$, we see that $\chi(G - v) \leq \Delta$ for all $v \in V(G)$. In particular, G is Δ -regular.

First, suppose G is 3-regular. If G contains a diamond D , then we may 3-color $G - D$ and easily extend the coloring to D by first coloring the nonadjacent vertices in D the same. So, G doesn't contain diamonds. Since G is not a forest it contains an induced cycle C . Since $K_4 \not\subseteq G$ we have $|N(C)| \geq 2$. So, we may take different $x, y \in N(C)$ and put $H := G - C$ if x is adjacent to y and $H := (G - C) + xy$ otherwise. Then, H doesn't contain K_4 as G doesn't contain diamonds. By minimality of $|G|$, H is 3-colorable. That is, we have a 3-coloring of $G - C$ where x and y receive different colors. We can easily extend this partial coloring to all of G since each vertex of C has a set of two available colors and some pair of vertices in C get different sets.

Hence we must have $\Delta \geq 4$. Consider a Δ -coloring of $G - v$ for some $v \in V(G)$. Each color must be used on every K_Δ in $G - v$ and hence some color must be used on every K_Δ in G . Let M be such a color class expanded to a maximal independent set. Then $\chi(G - M) = \chi(G) - 1 = \Delta > \max\{3, \omega(G - M), \Delta(G - M)\}$, a contradiction. \square

We note that the reduction to the cubic case is an immediate consequence of more general lemmas on hitting all maximum cliques with an independent set (see [5], [8] and [4]). H. Tverberg pointed out that this reduction was also demonstrated in his paper [9].

LIFTING TO (ONLINE) LIST COLORING

In [6], Kostochka and Yancey gave a simple, yet powerful, application of the Kernel Lemma to a list coloring problem. In [3], we generalized their idea to get a widely applicable general tool. As an illustration, we use a special case of this tool to prove the list coloring version of Brooks' Theorem (first proved by Vizing [10]). The same argument proves the online list coloring version of Brooks' Theorem, but we will stick to ordinary list coloring for simplicity.

A *kernel* in a digraph D is an independent set $I \subseteq V(D)$ such that each vertex in $V(D) - I$ has an edge into I . A digraph in which every induced subdigraph has a kernel is called *kernel-perfect*. A simple induction argument shows that kernel-perfect orientations can be very useful for list coloring:

Kernel Lemma. *Let G be a graph and $f: V(G) \rightarrow \mathbb{N}$. If G has a kernel-perfect orientation such that $f(v) \geq d^+(v) + 1$ for each $v \in V(G)$, then G is f -choosable.*

All bipartite graphs are kernel-perfect, the following lemma from [6] generalizes this fact.

Lemma 1. *Let A be an independent set in a graph G and put $B := V(G) - A$. Any digraph created from G by replacing each edge in $G[B]$ by a pair of opposite arcs and orienting the edges between A and B arbitrarily is kernel-perfect.*

Proof. In a minimum counterexample G , to get a contradiction it suffices to construct a kernel. But this is easy since either A is a kernel or there is some $v \in B$ which has no out neighbors in A . In the latter case, a kernel in $G - v - N(v)$ together with v is a kernel in G . \square

Theorem 2. *Every graph satisfies $\chi_l \leq \max\{3, \omega, \Delta\}$.*

Proof. Suppose the theorem is false and choose a counterexample G minimizing $|G|$. Put $\Delta := \Delta(G)$. Using minimality of $|G|$, we see that $\chi_l(G - v) \leq \Delta$ for all $v \in V(G)$. In particular, G is Δ -regular.

By Brooks' Theorem, $\chi(G) \leq \Delta$ and hence $\alpha(G) \geq \frac{|G|}{\Delta}$. Let A be a maximum independent set in G and put $B := V(G) - A$. Then there are $\alpha(G)\Delta \geq |G|$ edges between A and B . Hence there exist nonempty induced subgraphs H of G with at least $|H|$ edges between $A \cap V(H)$ and $B \cap V(H)$. Pick such an H minimizing $|H|$. For $v \in V(H)$, let d_v be the number of edges between $A \cap V(H)$ and $B \cap V(H)$ incident to v . We'll use minimality of $|H|$ to show that $d_v = 2$ for all $v \in V(H)$. Clearly, $d_v \geq 2$. Suppose $d_v < d_w$ for some other $w \in V(H)$. Then there are at least $|H| + d_w - 2$ edges across and hence removing v violates minimality. So, $d_v = d_w$ for all $v, w \in V(H)$. Call this common value d . Then there are at least $(d - 1)|H|$ edges across. Since $(d - 1)|H| - d \geq |H| - 1$ for $d > 2$, applying minimality shows that $d = 2$. Hence the edges between $A \cap V(H)$ and $B \cap V(H)$ induce a disjoint union of cycles.

Create a digraph Q from H by replacing each edge in $H[B \cap V(H)]$ by a pair of opposite arcs and orienting the edges between $A \cap V(H)$ and $B \cap V(H)$ around their respective cycles. Then Q is kernel-perfect by Lemma 1. Since $d^+(v) \leq d(v) - 1$ for each $v \in V(Q)$, applying the Kernel Lemma shows that H is f -choosable where $f(v) := d(v)$ for all $v \in V(H)$. But then given any Δ -list-assignment on G , we can color $G - H$ from its lists by minimality of $|G|$ and then complete the coloring to H , a contradiction. \square

The ad-hoc orientation construction in the above proof can be replaced with the following general lemma. This lemma follows easily by taking an arbitrary orientation and repeatedly reversing paths if doing so gets a gain (really, this is just the proof of the max-flow min-cut theorem). This can also be proved by splitting vertices and applying Hall's theorem.

Lemma 3. *Let G be a graph and $g: V(G) \rightarrow \mathbb{N}$. Then G has an orientation such that $d^-(v) \geq g(v)$ for all $v \in V(G)$ iff for every $H \trianglelefteq G$ we have*

$$\|H\| + \|H, G - H\| \geq \sum_{v \in V(H)} g(v).$$

Using Lemma 3 in a similar way to the proof of Brooks' Theorem for list coloring above, we get the general tool from [3].

Lemma 4. *Let G be a nonempty graph and $f: V(G) \rightarrow \mathbb{N}$ with $f(v) \leq d_G(v) + 1$ for all $v \in V(G)$. If there is independent $A \subseteq V(G)$ such that*

$$\|A, G - A\| \geq \sum_{v \in V(G)} d_G(v) + 1 - f(v),$$

then G has a nonempty induced subgraph H that is (online) f_H -choosable where $f_H(v) := f(v) + d_H(v) - d_G(v)$ for $v \in V(H)$.

In the case of Brooks' theorem, we want $f(v) := d_G(v)$, so the right side of the condition in Lemma 4 just equals $|G|$ and thus the existence of the desired A follows immediately from $\alpha(G) \geq \frac{|G|}{\Delta}$. More generally, as shown in [3], the classification of (online) degree-choosable graphs is a quick corollary of Lemma 4.

REFERENCES

- [1] R.L. Brooks, *On colouring the nodes of a network*, Mathematical Proceedings of the Cambridge Philosophical Society, vol. 37, Cambridge Univ Press, 1941, pp. 194–197.
- [2] R. Diestel, *Graph Theory*, Fourth ed., Springer Verlag, 2010.
- [3] H. Kierstead and L. Rabern, *Improved lower bounds on the number of edges in list critical and online list critical graphs*, In Preparation.
- [4] A.D. King, *Hitting all maximum cliques with a stable set using lopsided independent transversals*, Journal of Graph Theory **67** (2011), no. 4, 300–305.
- [5] A.V. Kostochka, *Degree, density, and chromatic number*, Metody Diskret. Anal. **35** (1980), 45–70 (in Russian).
- [6] A. Kostochka and M. Yancey, *Ore's Conjecture on color-critical graphs is almost true*, arXiv:1209.1050 (2012).
- [7] L. Lovász, *Three short proofs in graph theory*, Journal of Combinatorial Theory, Series B **19** (1975), no. 3, 269–271.
- [8] L. Rabern, *On hitting all maximum cliques with an independent set*, Journal of Graph Theory **66** (2011), no. 1, 32–37.
- [9] H. Tverberg, *On Brooks' theorem and some related results*, Mathematics Scandinavia **52** (1983), 37–40.
- [10] V.G. Vizing, *Vertex coloring with given colors*, Metody Diskretn. Anal. **29** (1976), 3–10 (in Russian).